



**UNIVERSITI PUTRA MALAYSIA**

**CYTOTOXIC EFFECTS OF METHYLGERAMBULLIN AND BIS  
(METHYLTHIOMETHYL)-DISULPHIDE (SB) ON T-LYMPHOBLASTIC  
LEUKEMIC CELL LINE (CEM-SS)**

**SHAR MARIAM MOHAMED**

**FSMB 2000 15**

**CYTOTOXIC EFFECTS OF METHYLGERAMBULLIN AND BIS-  
(METHYLTHIOMETHYL)-DISULPHIDE (SB) ON T-LYMPHOBLASTIC  
LEUKEMIC CELL LINE (CEM-SS)**

**By**

**SHAR MARIAM MOHAMED**

**Thesis Submitted in Fulfilment of the Requirements for the  
Degree of Doctor of Philosophy in the Faculty of  
Food Science and Biotechnology  
Universiti Putra Malaysia**

**August 2000**



## DEDICATION

Dedicated to my dear husband, I cannot thank you enough for your love and support. Having the blessings of being your wife is certainly my wildest dream come true. Your incredibly unique love has touched me in ways I cannot describe and I am deeply thankful (syukur) to the Almighty that I've been given the chance to weave my love with yours. My deepest gratitude and love to my dear parents Mohamed Hj. Ismail and Hasnah Midon, sister Ayu and brother Don for their everlasting eternal support, courage and love, which keep me going as a stronger person each day. I am always with you more than ever. To my in-laws, I am so blessed with your warmth and support..... thank you.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirements for the degree of Doctor of Philosophy

**CYTOTOXIC EFFECTS OF METHYLGERAMBULLIN AND BIS-  
(METHYLTHIOMETHYL)-DISULPHIDE (SB) ON T-LYMPHOBLASTIC  
LEUKEMIC CELL LINE (CEM-SS)**

**By**

**SHAR MARIAM MOHAMED**

**August 2000**

**Chairman: Associate Professor Abdul Manaf Ali, Ph.D.**

**Faculty: Food Science and Biotechnology**

The cytotoxic effects of 2 sulphur-containing compounds were studied on T-lymphoblastic leukemic cell line. Methylgerambullin is believed to be a new sulphone derived from a methylthiopropenoic acid isolated from *Glycosmis calcicola* (family *Rutaceae*). Another sulphonic compound is bis-(methylthiomethyl)-disulphide, an extract from *Scorodocarpus borneensis* (family *Olacaceae*) with irritating garlic-like odor. Cytotoxic activities of methylgerambullin and bis-(methylthiomethyl)-disulphide were tested against CEM-SS (T-lymphoblastic leukaemia), KU812F (chronic myelogeneous leukaemia), UACC-62 (melanoma) and HT29 (colon cancer) cell lines using MTT, a colorimetric tetrazolium-based assay. Cytotoxic concentrations of the compounds that killed cells by 50% (CD50) with respect to untreated cell population, varied among the cell lines tested. CEM-SS was found to be the most sensitive cell line to methylgerambullin and bis-(methylthiomethyl)-disulphide with  $CD50 = 0.25 \mu\text{g/ml}$  and  $3.50 \mu\text{g/ml}$  respectively. The cytotoxic effects exerted by both compounds on this cell line was studied from both morphological

manner over 72 hours period. Microscopic observations, including inverted microscopy of live cultures, fluorescent microscopy of acridine orange-propidium iodide stained cultures, and scanning and transmission electron microscopy showed that both necrotic and apoptotic death occurred in methylgerambullin- and bis-(methylthiomethyl)-disulphide-treated cell populations, based on morphological criteria. From agarose gel electrophoresis and quantitative analyses of internucleosomal cleavage, treatments with these compounds at their respective CD50 doses did not yield random or multiple of 180-200 bp DNA fragmentation which often associated with necrotic and apoptotic deaths respectively. Such observation may simply owe to the fact that the percentage of apoptosis and necrosis events were fairly low as quantified after acridine orange-propidium iodide staining, or may also suggest the involvement of sulphur residue in methylgerambullin and bis-(methylthiomethyl)-disulphide which act as an antioxidant, thus protecting DNA degradation from occurring. Flow cytometric analyses based on annexin V-FITC (fluorescein isothiocyanate) binding to the phosphatidylserines residue which was translocated from the inner to the outer leaflet of the plasma membrane showed that the onset of apoptosis in both methylgerambullin- and bis-(methylthiomethyl)-disulphide-treated population was at 6 hours exposure. Both methylgerambullin and bis-(methylthiomethyl)-disulphide induced G0/G1 arrest up to 48 hours and 24 hours respectively followed by arrest in the subsequent S phase.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah.

**KESAN SITOTOKSIK OLEH METHYLGERAMBULLIN AND BIS-(METHYLTHIOMETHYL)-DISULPHIDE (SB) KE ATAS SEL-SEL T-LIMFOBLASTIK LEUKEMIA (CEM-SS)**

**Oleh**

**SHAR MARIAM MOHAMED**

**Ogos 2000**

**Pengerusi: Profesor Madya Dr. Abdul Manaf Ali**

**Fakulti: Sains Makanan dan Bioteknologi**

Kesan sitotoksik oleh 2 sebatian yang mengandungi sulfur telah dikaji ke atas sel-sel T-limfoblastik leukemia. Metilgerambullin, dipercayai satu sebatian sulfon yang terbaru ditemui adalah hasil terbitan daripada asid metiltiopropenoik yang diekstrak daripada *Glycosmis calcicola* (keluarga *Rutaceae*). Satu lagi sebatian ialah bis-(metiltiometil)-disulfida yang diekstrak daripada *Scorodocarpus borneensis* (keluarga *Olacaceae*) yang berbau seperti bawang putih. Aktiviti sitotoksik metilgerambullin dan bis-(metiltiometil)-disulfida telah diuji ke atas sel-sel CEM-SS (T-limfoblastik leukemia), KU812F (kronik myelogeneous leukemia), UACC-62 (melanoma) dan HT29 (kanser kolon) menggunakan kaedah MTT yang bergantung kepada perubahan warna tetrazolium. Hasil kajian menunjukkan bahawa sel CEM-SS adalah sel yang paling sensitif terhadap metilgerambullin dan bis-(metiltiometil)-disulfida dengan kepekatan yang berupaya membunuh 50% daripada populasi sel (CD50) iaitu 0.25 µg/ml bagi metilgerambullin dan 3.50 µg/ml bagi bis-(metiltiometil)-disulfida. Kesan sitotoksik sebatian-sebatian ini dinilai melalui perubahan pada morfologi sel dan

perubahan di peringkat molekular selepas rawatan dengan sebatian-sebatian tersebut selama 72 jam. Penilaian mikroskopik termasuklah menggunakan mikroskop terbalikan ke atas kultura hidup, mikroskop floresen selepas pewarnaan sel dengan akridin oren- propidium iodida dan mikroskopi elektron menunjukkan bahawa kematian apoptosis dan nekrosis berlaku di dalam populasi sel yang dirawat dengan metilgerambullin dan bis-(metiltiometyl)-disulfida. Daripada keputusan yang diperolehi melalui elektroforesis pada jel agaros dan analisis kuantitatif pada belahan nukleosom, rawatan dengan kedua-dua sebatian pada dos CD50 masing-masing tidak menghasilkan belahan rawak atau belahan spesifik 180-200 bp pada DNA yang masing-masing dikaitkan dengan kematian nekrosis dan apoptosis. Pemerhatian tersebut mungkin disebabkan oleh peratusan kematian apoptosis dan nekrosis yang terlalu sedikit sepertimana yang telah dikenalpasti secara kuantitatif melalui kaedah pewarnaan sel dengan akridin oren dan propidium iodida, atau mungkin disebabkan oleh kesan perlindungan sulfur yang terdapat pada methylgerambullin dan bis-(metiltiometyl)-disulfida yang bertindak sebagai antioksidan yang melindungi daripada berlakunya belahan DNA. Analisis menggunakan flow saktometer berdasarkan pada ikatan annexin V-FITC (fluoresen isotiosianida) pada fosfatidilserine yang mengalami translokasi dari lapisan dalam ke lapisan luar membran plasma, menunjukkan bahawa kejadian apoptosis dalam populasi sel yang dirawat dengan methylgerambullin dan SB berlaku seawal 6 jam selepas rawatan. Methylgerambullin dan SB juga didapati berupaya merencatkan sel pada fasa G0/G1 dalam kitaran sel hingga 48 jam dan 24 jam masing-masing selepas rawatan, diikuti dengan rencatan pada fasa S dalam kitaran sel.

## **ACKNOWLEDGEMENTS**

I would like to especially thank my supervisor, Associate Professor Dr. Abdul Manaf Ali for his help, guidance, encouragement, criticism and patience, without whom I would not be able to go through my research years successfully. Above all, thank you so much for being so generous with ideas. My utmost appreciation to my supervisory committee members, Associate Professor Dr. Mawardi Rahmani, Dr. Jasbir S. Dhaliwal and Dr. Raha Abdul Rahim for being very helpful throughout my research years.

For kindly allowing me to use all the facilities within the Faculty of Food Science and Biotechnology, I thank the Dean, Professor Dr. Gulam Rasul Rahmat Ali. To all the staffs of the faculty, thank you so much for the technical assistance given. Not forgetting the Universiti Putra Malaysia, I am truly grateful for the financial support from which I was able to complete my research work successfully.

I gratefully acknowledge the following people for their time, advice and generous contributions for making this piece of work possible: staffs of the Electron Microscopy Unit, Faculty of Veterinary Medicine; staffs of the Immunology Division, Institute for Medical Research Kuala Lumpur especially Mr. Ong, Mr. Quek, Kak Zaitun and Kak Zuraidah; Dr. Leong Kah Ho from Becton Dickinson; and last but not least the staffs of Makmal Penyelidikan Pusat such as Kak Fadillah and Kak Liza. To my fellow Animal Cell Culture Lab colleagues; Tony, Lim, Khor, Kak Niza, Kak Siti, Yih Yih, and all my juniors, thanks a zillion for sharing the work load of keeping the lab tidy with me (you know I got a little



upset once in a while when the contaminants were not discarded properly). To have the wonderful opportunity to work with you guys and to experience the enjoyable moments is an everlasting souvenir that I will always cherish.

I certify that an Examination Committee met on 28 August 2000 to conduct the final examination of Shar Mariam Mohamed on her Doctor of Philosophy thesis entitled "Cytotoxic Effects of Methylgerambullin and Bis-(methylthiomethyl)-disulphide (SB) on T-Lymphoblastic Leukemic Cell Line (CEM-SS)" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Professor Suhaila Mohamed, Ph.D,  
Faculty of Food Science and Biotechnology,  
Universiti Putra Malaysia  
(Chairperson)


Associate Professor Abdul Manaf Ali, Ph.D,  
Faculty of Food Science and Biotechnology,  
Universiti Putra Malaysia  
(Member)

Associate Professor Mawardi Rahmani, Ph.D,  
Faculty of Environmental Science,  
Universiti Putra Malaysia  
(Member)

Raha Abdul Rahim, Ph.D,  
Faculty of Food Science and Biotechnology,  
Universiti Putra Malaysia  
(Member)

Jasbir S. Dhaliwal, Ph.D,  
Division of Immunology,  
Institute for Medical Research, Kuala Lumpur  
(Member)

Professor Peter J. Houghton, Ph.D,  
King's College of London  
(Independent Examiner)



MOHD. GHAZALI MOHA YIDIN, Ph.D.  
Professor/Deputy Dean of Graduate School,  
Universiti Putra Malaysia

Date: 01 NOV 2000

This thesis submitted to the Senate of Universiti Putra Malaysia and was accepted as fulfilment of the requirements for the degree of Doctor of Philosophy.



---

KAMIS AWANG, Ph.D,  
Associate Professor  
Dean of Graduate School,  
Universiti Putra Malaysia

Date: **14 DEC 2000**

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



---

Candidate  
SHAR MARIAM MOHAMED

Date: 25 October 2000

## TABLE OF CONTENTS

	Page
<b>DEDICATION .....</b>	<b>ii</b>
<b>ABSTRACT.....</b>	<b>iii</b>
<b>ABSTRAK.....</b>	<b>v</b>
<b>ACKNOWLEDGEMENTS .....</b>	<b>vii</b>
<b>APPROVAL SHEETS.....</b>	<b>ix</b>
<b>DECLARATION .....</b>	<b>xi</b>
<b>LIST OF TABLES.....</b>	<b>xv</b>
<b>LIST OF FIGURES.....</b>	<b>xvi</b>
<b>LIST OF PLATES .....</b>	<b>xviii</b>
<b>LIST OF ABBREVIATIONS.....</b>	<b>xix</b>
<b>CHAPTER</b>	
<b>I INTRODUCTION .....</b>	<b>1</b>
<b>II LITERATURE REVIEW .....</b>	<b>6</b>
Sulphur-containing Compounds .....	6
Leukemia.....	12
Cell Death	
Nomenclature and Terminology .....	15
Biochemical and Genetic Control of Cell Death .....	22
Mitochondria As Damage Sensor and Execution of Cell Death....	23
Nuclear Events in Apoptosis.....	27
Cell Death and Its Relevance in Animal Development and	
Disease .....	31
Apoptosis in Hematopoiesis and Hematological Malignancies....	33
Cell Death Detection	
Screening for potential anticancer drugs.....	36
Microculture Tetrazolium Assay (MTA) .....	36
Microscopic Analyses .....	40
Basic Principle of Flow Cytometry .....	44
The Cell Cycle .....	45
Flow Cytometric Analyses of DNA Content in Human Malignancy	50
Cell Cycle and Its Significance in Cytotoxicity and Chemotherapy ..	52
Detection of Apoptosis Using Flow Cytometric Method .....	55
<b>III MATERIALS AND METHODS.....</b>	<b>59</b>
Cell Line .....	59
Reviving Cells.....	59
Storing Cells .....	60
Cell Line Maintenance	
Suspension Cell Culture .....	60



	Anchorage-dependent Cell Culture.....	60
	Isolation of Human Peripheral Blood Mononuclear Cells.....	61
	Test Compounds	
	Purified Natural Products.....	61
	Commercial Drugs .....	63
	Cytotoxic Assay .....	64
	Statistical Analysis.....	65
	Proliferation Assay.....	65
	Synchronization of Cells.....	66
	Morphological Analyses	
	Giemsa Staining.....	66
	Acridine Orange (AO) and Propidium Iodide (PI) Staining.....	67
	Electron Microscopy .....	67
	Molecular Analyses	
	DNA Extraction and Analyses for Apoptosis Induction.....	69
	Cell Death Detection ELISA <sup>PLUS</sup> kit.....	70
	Flow Cytometric Analyses	
	Cell Cycle Specificity .....	71
	Early Apoptosis Detection (Annexin V binding).....	72
IV	RESULTS .....	73
	Cytotoxicity of Test Compounds Against Various Cell Lines .....	73
	Statistical Analyses .....	78
	Proliferative Profiles of CEM-SS After Treatment With Various Test Compounds at Their Respective CD50 .....	82
	Cellular Examination	
	Determination of CD50 from MTT Assay: Dose-dependent Studies.....	84
	Treatment at CD50: Time-dependent Studies .....	91
	Effects of Control Drugs on CEM-SS .....	94
	Effects of Methylgerambullin and SB on Peripheral Blood Mononuclear Cells.....	96
	Fluorescence Microscopy .....	96
	Giemsa Staining.....	105
	Electron Microscopy .....	105
	Progression of Apoptosis Events .....	116
	Progression of Necrosis Events .....	119
	Synchronization of Cells.....	119
	Effects of Methylgerambullin and SB on Nuclear Fragmentation.....	121
	Flow Cytometric Analyses	
	Detection of Early Apoptosis Events.....	125
	Cell Cycle Analyses .....	135
V	DISCUSSION .....	152
	Cytotoxic, Proliferative and Resistance Profiles of Test Compounds .....	152

Proliferative Profiles of CEM-SS After Treatment With Test Compounds .....	156
Effects of Methylgerambullin and SB on Cell Morphology .....	162
Effects of Methylgerambullin and SB on Nuclear Fragmentation.....	173
Flow Cytometry: Detection of Apoptosis .....	182
Flow Cytometry: Cell Cycle Analyses.....	186
VI     CONCLUSION.....	194
<b>REFERENCES.....</b>	<b>200</b>
<b>APPENDICES.....</b>	<b>219</b>
A     Preparation of Media and Solutions.....	220
B     Calculation for Cell Viability and Cell Concentration .....	222
C     The Means and Standard Error of Means (SEM) Corresponding to Figures 12 and 13.....	223
D     Two-ways Analysis of Variance (ANOVA).....	224
<b>VITA . .....</b>	<b>228</b>

## LIST OF TABLES

Table	Page
1 ..... Biochemical determinants of apoptosis and necrosis.....	26
2 ..... Commercial drugs of various classes used as controls in this study.	63
3 ..... Sensitivity of various cell lines to methylgerambullin and SB .....	73
4 ..... Sensitivity of CEM-SS to various commercial drugs .....	78
5 ..... Sensitivity of CEM-SS to methylgerambullin, SB and various commercial drugs .....	80
6 ..... The percentage of mitotic cells induced by colchicine at different concentrations.....	121
7 ..... The summary of dose-dependent percentage of viable, apoptotic and necrotic CEM-SS cells .....	129
8 ..... The summary of time-dependent percentage of viable, apoptotic and necrotic CEM-SS cells .....	134
9 ..... Cell cycle distribution in CEM-SS after 24 hours incubation with methylgerambullin, SB, gonoiothalamine, cisplatin, etoposide and colchicine at various concentrations.....	144
10 ..... Cell cycle distribution in CEM-SS after incubation with methylgerambullin, SB, gonoiothalamine, cisplatin, etoposide and colchicine at CD50 doses for various duration of time.....	150
11 ..... Time-dependent cytotoxic profiles of control drugs, methylgerambullin and SB against CEM-SS cell line.....	223
12 ..... Proliferative profiles of CEM-SS cells after treatment with methylgerambullin, SB and control drugs at their respective CD50 dose.....	223
13 ..... The summary of means of CD50 for different compounds/drugs (Xi) and means of CD50 exhibited by different CEM-SS sub-populations at different time (Xj) .....	224
14 ..... The summary of two-ways ANOVA.....	225





## LIST OF FIGURES

Figure	Page
1 ..... The experimental approach implemented in the current study .....	5
2 ..... The structure of methylgerambullin and SB (bis-(methylthiomethyl)-disulphide).....	11
3 ..... Morphological changes during apoptosis and necrosis.....	20
4 ..... Flow diagrams of apoptotic and necrotic major events.....	21
5 ..... Tetrazolium salts and its formazan.....	37
6 ..... Evolution of tetrazolium salts used in measuring cell viability .....	38
7 ..... The cell cycle .....	48
8 ..... Cyclical changes in DNA content of a dividing cell.....	49
9 ..... The percentage of cell viability of various cell lines after treatment with different concentrations of methylgerambullin for 72 hours .....	75
10 ..... The percentage of cell viability of various cell lines after treatment with different concentrations of SB for 72 hours .....	76
11 ..... The percentage of cell viability of CEM-SS after treatment with different concentrations of control drugs, methylgerambullin and SB for 72 hours <sup>74</sup> .....	77
12 ..... Time-dependent cytotoxic profiles of control drugs, methylgerambullin and SB against CEM-SS cell line.....	81
13 ..... Proliferative profiles of CEM-SS cell line .....	83
14 ..... Time-dependent trends of apoptotic and necrotic incidence in CEM-SS cell population.....	103
15 ..... Detection of mono- or oligonucleosomal DNA fragmentation in methylgerambullin- and SB-treated CEM-SS population by ELISA <sup>PLUS</sup> immunoassay .....	124
16 ..... Classification of viable, apoptosis and necrosis based on Annexin V-FITC and propidium iodide uptake.....	125
17 ..... Flow cytometric analyses on detection of apoptosis and necrosis at different doses for 24 hours.....	127

18 .....	Time course flow cytometric analyses on detection of apoptosis and necrosis at CD50 doses .....	132
19 .....	Determination of cell cycle statistics from dot plot and histogram using CELLFIT and LYSIS II softwares .....	138
20 .....	Flow cytometric cell cycle analyses based on DNA content .....	142
21 .....	Time-dependent flow cytometric cell cycle analyses based on DNA content .....	148
22 .....	Possible stages in the process of responding to cellular damage .....	153
23 .....	Characterization of viable, apoptotic and necrotic cells based on the chromatin structure after dual staining with acridine orange and propidium iodide .....	167
24 .....	The distribution of means between samples in the hypothesis testing .....	226
25 .....	The distribution of means within samples in the hypothesis testing .....	226

## LIST OF PLATES

Plate	Page
1 ..... Photomicrographs from inverted microscopy .....	85
2 ..... Photomicrographs of cells after the addition of MTT .....	89
3 ..... Photomicrograph of formazan crystals surrounding viable CEM-SS cells detected after 4 hours incubation with MTT .....	90
4 ..... Time-dependent morphological changes of CEM-SS cells treated with methylgerambullin .....	92
5 ..... Time-dependent morphological changes of CEM-SS cells treated with SB .....	93
6 ..... CEM-SS cells after 72 hours treatment with control drugs.....	95
7 ..... Peripheral blood mononuclear cells .....	97
8 ..... Photomicrographs of acridine-orange-propidium iodide stained CEM-SS cells.....	100
9 ..... Photomicrographs of viable, apoptotic and necrotic cells.....	102
10 ..... Giemsa staining .....	106
11 ..... Scanning electron micrographs of methylgerambullin-treated CEM-SS cells.....	109
12 ..... Scanning electron micrographs of SB-treated CEM-SS cells.....	111
13 ..... Close up view of scanning electron micrographs.....	113
14 ..... Transmission electron micrographs of CEM-SS treated with methylgerambullin and SB.....	115
15 ..... Ultrastructural changes in apoptotic cells .....	117
16 ..... Ultrastructural changes in necrotic cells .....	120
17 ..... DNA profiles from agarose gel electrophoresis.....	123

## **LIST OF ABBREVIATIONS**

2-CDA .....	chloro deoxyadenosine
ABTS .....	2,2'-Azino-di[3-ethylbenzthiazolin-sulfonat]
ADP ... ..	adenosine diphosphate
AIF.....	apoptosis inducing factor
ALL ... ..	acute lymphoblastic leukemia
AML .. .....	acute myeloid leukemia
AO .....	acridine orange
Ara-c .. .....	arabinofuranosylcytosine
ATP ... .....	adenosine triphosphate
CA .....	calyculin A
CD50 . .....	cytotoxic dose resulting in 50% reduction of cell population
CGM.. .....	complete growth medium
CLL.....	chronic lymphocytic leukemia
CML .. .....	chronic myeloid leukemia
CO <sup>2</sup> .....	carbon dioxide
CV .....	coefficient of variation
Cyt-c .. .....	cytochrome c
DCF ... .....	deoxycorformycin
DDS.....	diaminodiphenyl sulphone
DMSO .....	dimethyl sulfoxide
DNA .. .....	deoxyribonucleic acid
ER.....	endoplasmic reticulum
FDA ... .....	Food and Drug Administration
FITC .. .....	fluorescein isothiocyanate
FSC.....	forward scatter
GSH.....	glutathione
HCl .....	hydrochloric acid
HIV.....	human immunodeficiency virus
ICAD . .....	inhibitor of caspase-activated DNase
ICE.....	interleukin converting enzyme
KCl .....	potassium chloride
MTT .. .....	3(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
MTA .. .....	microculture tetrazolium assay
MTS.....	3-[4,5-dimethylthiazol-2-yl]-5-(carboxymethoxyphenyl),(4-sulphophenyl)-2H-tetrazolium
MeOH .....	methanol
MTP.....	microtiter plate
NaCl .. .....	sodium chloride
NHAC .....	N4-hexadecyl-1-β-D-arabinofuranosylcytosine
OD .....	optical density
OA.....	okadaic acid
PARP . .....	poly(ADP-ribose) polymerase
PBMC .....	peripheral blood mononuclear cell
PBS.....	phosphate buffered saline
PI .....	propidium iodide
PMS.....	phenazine methosulphate



PMT ...	photo multiplier
PS.....	phosphatidylserine
PT .....	permeability transition
RNA....	ribonucleic acid
RNP ...	ribonucleoprotein particles
ROS ...	reactive oxygen species
SB .....	bis-(methylthiomethyl)-disulphide isolated from <i>Scorodocarpus borneensis</i> Becc.
SDS....	sodium dodecyl sulphate
SEM....	scanning electron microscope
SSC....	side scatter
TEM ..	transmission electron microscope
TNF ...	tumor necrosis factor
TUNEL.....	Tdt-mediated dUTP-biotin nick end labelling
UV .....	ultraviolet
XTT ...	sodium (2,3-bis[2-methoxy-4-nitro-5-sulphophenyl]-2H-tetrazolium-5-carboxanilide)

## **CHAPTER I**

### **INTRODUCTION**

Screening for potential anticancer compounds has been one of the most important aspects in cancer research. Many commercially available drugs are synthetically derived, for example etoposide, a semi-synthetic epipodophyllotoxin that is one of the most commonly used anti-cancer drugs (Joel, 1996). Although these synthetic drugs are known to be very effective against cancer cells, they also possess some undesirable side effects such as neurotoxicity, fibrinolysis and ototoxicity (hearing loss) (Casciato and Lowitz, 1995). Thus, choices for improvement are confined to introduce natural products as potential anticancer agents. Phytochemicals, one of these natural substances have been known to exhibit a wide variety of medicinal value ever since the ancient times. The scientific theories underlying their medicinal properties however remained unknown then. Wilkinson, (1998) recently highlighted that herbal products have great potential in the emerging nutraceutical and pharmaceutical industries in that they are widely consumed as food and are used in preventive and curative treatments throughout the world. It was reported in the same study that tropical forests have produced 47 major pharmaceutical drugs of world-wide importance and it is estimated that 328 potential drugs of major importance is yet to be discovered which would be worth \$147 billion. The tropical forests still offers 125,000 flowering plant species that are of pharmacological relevance but this will involve 50,000 to one million screening

tests to discover one profitable drug, which seem to be the ultimate constraint in the discovery of new drugs.

It has been reported by the Alliance Pharmaceutical Corporation that on average, 12 years are needed for an experimental drug to be brought from the laboratory to the market shelf (Wierenga and Eaton, Internet). Preclinical testing requires three and a half years of experiments in laboratory on *in vitro* and animal models to assess safety and biological activities of the compound. Another six years are required for clinical trials that assess the safety, dosage, effectiveness and monitor the adverse reaction of the compound. Only five in 5000 compounds that enter preclinical testing make it to human testing. Approval process by the Food and Drug Administration (FDA) will take about two and a half years where normally only one of the five compounds tested in humans is approved.

Despite the duration of time required to evaluate a potential drug, numerous phytochemicals have been scientifically isolated and their discovery was very much directed in the perspective of anticancer drug development. Plant constituents such as taxol, camptothecin, podophyllotoxin, vinblastine and vincristine represent some of the most important drugs currently utilized for the treatment of human cancers (Wall and Wani, 1993). Numerous more phytochemicals are under investigations and have the potential to be on the pharmaceutical market shelves. Among these are betulinic acid, gonoithalamin and dammarane-type triterpenes, to name a few. Betulinic acid, which is available in abundant supply from the bark of white birch trees was reported to

be specific for inhibiting human melanomas (Pisha *et al.*, 1995). Gonoiothalamine, isolated from various *Gonoiothalamus* sp. was found to be toxic to cervical, pancreas, gastric and breast carcinoma (Ali *et al.*, 1996). Dammarane-type triterpenes from *Cloeme africana* has been shown to be highly toxic against leukemia (Nagaya *et al.*, 1997).

One of the most extensively studied diseases is leukemia that remains to be a formidable disease. Due to the ease of obtaining repeated samples from blood, marrow or lymph nodes, most of the principles of cancer therapy is largely based on the leukemia model. Important breakthroughs in cancer therapy were achieved in the treatment of leukemia, which includes the use of combination therapy, immunotherapy, supportive therapy using antibiotics and blood product support and bone marrow transplant (Ng, 1996). Treatment with antileukaemic drugs however only controls the symptom without enhancing the survival rate (Arthur, 1989). Chemotherapeutic agents, also known as cytotoxins due to its ability to kill cells are normally less specific in that these agents kill not only cancer cell but also normal cells. Thus, the most apparent implication of chemotherapy is emanation of side effects which include hair loss, nausea, vomiting, constipation and can be as disastrous as myelosuppression (drop in blood count – neutropenia, thrombocytopenia, anaemia), and damage on cardiac, pulmonary, renal and hepatic systems (Ng, 1996). This prompted efforts to screen for antileukaemic agents that will have a dramatic cytotoxic effect specifically on the target cells. Most of the drugs currently used act via two distinct mechanisms: inhibition of cell proliferation or induction of active cell



death (Vial *et al.*, 1997). Compounds that act via the latter mechanism are likely to be a better candidate as an antileukaemic agent because in the former action, some individual cells may have escaped through salvage pathways, and will continue to proliferate. The advancement in research techniques and the introduction of new molecular tools in recent years have prompted more efforts in investigations to build understanding on how exposure to the existing or newly discovered drugs lead to cell death. The efficacy of these drugs has been critically assessed by the many mechanisms of cell death and resistance that adversely limit their efficacy. Today, due to the urgent need for studies in various fields that closely mimics the animal systems, animal cell culture which was only an esoteric art during those time has become an established technology. New drug discovery, for example, may not be recognized and may be unsafe for use in man. Thus, pre-clinical trials are needed, which can be made possible with cell culture technology.

It is thus crucial to determine the mode of action exerted by a potential anticancer agent. This has led to the design of the experimental approach which will be implemented in the current study. Both cellular and molecular changes in treated cell populations will be examined and compared with control (untreated) populations in a time course manner. Hence, either necrotic or apoptotic cell death could be suggested. Morphological studies may offer answers to whether physiological and accidental deaths are causally related and to the degree of overlapping between apoptotic and necrotic pathways.